

← **DOSING**
See back panel

durvet

DuraFend™

Safe-Guard® (fenbendazole) 0.5%

Type C Medicated Feed

**Dewormer for Beef & Dairy Cattle, Horses,
Zoo and Wildlife Animals**

**SAFE-GUARD® (fenbendazole) 0.5%
Medicated Dewormer Pellets**

ACTIVE DRUG INGREDIENT

fenbendazole.....0.5% (2.27 g/lb)

INGREDIENTS

Processed grain by-products, calcium carbonate, molasses products, roughage products, silicon dioxide, propionic acid (preservative), acetic acid, natural and artificial flavoring components, sodium propionate (preservative), potassium sorbate (preservative), propylene glycol and citric acid (preservative).

GUARANTEED ANALYSIS

Crude Protein, min.....11.5%
Crude Fat, min.....2.5%
Crude Fiber, max.....10%

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Safe-Guard is property of Intervet International B.V. or affiliated companies or licensors and are protected by copyrights, trademark and other intellectual property laws. ©2010 Intervet International B.V.

All rights reserved. • intervetusa.com • 800.521.5767

See back panel for instructions for use and warning statements.

MEASURING



Scoop Included
8 oz (0.23 KG)

NET WT: 5 LB (2.27 KG)



durvet

DuraFend™

Safe-Guard® (fenbendazole) 0.5%

CATTLE – Dairy and Beef: For the removal and

control of: Lungworms: (*Dictyocaulus viviparus*); Stomach worms: barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*); Intestinal worms: hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata*, *Cooperia oncophora*); Bankrupt worms: (*Trichostrongylus colubriformis*); and Nodular worms: (*Oesophagostomum radiatum*). **HORSES: For the control of:** Large strongyles: (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*, *Triodontophorus* spp.); Small strongyles: (*Trichostrongylus* spp., *Cylicocyclus* spp., *Cylicostephanus* spp.); Pinworms: (*Oxyuris equi*); and Ascarids: (*Parascaris equorum*).

ZOO AND WILDLIFE ANIMALS – Bighorn Sheep:

Lung worms: (*Protostrongylus* spp.). **Feral Swine:** Kidney worms: (*Stephanurus dentatus*); Roundworms (*Ascaris suum*); and Nodular worms (*Oesophagostomum dentatum*).

Ruminants (Subfamilies antilopinae, hippotraginae, and caprinae): Small stomach worm: (*Trichostrongylus* spp.); Thread-necked intestinal worm: (*Nematodirus* spp.); Barberpole worm (*Haemonchus* spp.); and Whipworm: (*Trichuris* spp.).

Contact Durvet, Inc. for other dewormer products that are cleared for use in Swine and Turkeys.

FEEDING DIRECTIONS – Enclosed scoop measures 8 oz.

CATTLE – DAIRY AND BEEF: 5 mg fenbendazole per kg body weight in a ONE (1) DAY treatment (2.27 mg fenbendazole per pound of bodyweight). Example: 1000 lb. cattle should receive 1 lb. of DuraFend 0.5% in one (1) day. Feed as the sole ration or top dress for one day. No prior withdrawal of feed or water necessary. Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks.

HORSES: For large strongyles, small strongyles and pinworms, the recommended dose is 5 mg fenbendazole per kg body weight (2.27 mg fenbendazole per pound) in a ONE (1) DAY treatment. Example: a 1000-lb horse should receive 1.0 pounds of DuraFend 0.5% in a ONE (1) DAY treatment. For ascarids, the recommended dose is 10 mg fenbendazole per kg body weight (4.54 mg fenbendazole per pound) in a ONE (1) DAY treatment. Example: a 1000-lb horse should receive 2.0 pounds of DuraFend 0.5% in a ONE (1) DAY treatment. All horses must be eating normally to ensure that each animal consumes an adequate amount of medicated feed. Regular deworming at intervals of 6 to 8 weeks may be

required due to the possibility of reinfection. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

ZOO AND WILDLIFE ANIMALS – Big Horn Sheep:

10 mg fenbendazole per kg of body weight per day, times (x) 3 days of treatment. **Feral Swine:** 3 mg fenbendazole per kg of body weight per day, times (x) 3 days of treatment.

Ruminants (subfamilies antilopinae, hippotraginae and caprinae):

2.5 mg fenbendazole per kg of body weight per day, times (x) 3 days of treatment. Example: 250 lb Feral Swine should receive 0.15 lbs. of DuraFend 0.5% per day for 3 days. It is recommended that the user exercise judgment expertise as needed for retreatment within six (6) weeks. This would depend upon the conditions of treated animals and ambient temperatures. Once correct proportions of Durafend 0.5% and feed ingredients have been established and blended, use as complete feed.

WARNINGS: KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. NOT FOR USE IN HUMANS. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, obtain an MSDS or for assistance, contact customer service at 1-800-441-8272.

Dairy and Beef – Residue warning – Cattle must not be slaughtered within 13 days following last treatment. There are no known contradictions to the use of the drug in cattle. For dairy cattle, there is no milk withdrawal period. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. **Horses:** Do not use in horses intended for human consumption. **Zoo and Wildlife Animals** - Residue warning – Do not use 14 days before or during the hunting season. Animal Safety: No contradictions for the use of fenbendazole in a zoo environment have been established. Administration to breeding and pregnant ruminants at 2 to 22 times the recommended dose has had no apparent adverse effect.



Manufactured for Durvet, Inc., Blue Springs, MO 64014
www.durvet.com 1-800-821-5570

NET WT: 5 LB (2.27 KG)

ISS 0916