

# Ivermectin Sheep Drench (ivermectin)

NDC 30798-637-12

0.08% Solution  
Parasiticide

## Description:

Ivermectin Sheep Drench is a ready-to-use, free-flowing solution of ivermectin. It is formulated to deliver the recommended dose rate of 0.2 mg ivermectin per 1 kg body weight given orally at a volume of 3.0 mL per 26 lbs body weight.

## Indications:

Ivermectin Sheep Drench provides treatment and control of adult and fourth-stage larvae of the following parasites: Gastrointestinal Roundworms – *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia curticei*, *Nematodirus spathiger*, *N. battus*, and *Oesophagostomum columbianum*; Lungworms – *Dictyocaulus filaria*; and all the larval stages of Nasal Bot – *Oestrus ovis*. It also provides treatment and control of adult forms only of the following Gastrointestinal Roundworms – *Haemonchus placei*, *Cooperia oncophora*, *Strongyloides papillosus*, *Oesophagostomum venulosum*, *Trichuris ovis*, and *Chabertia ovina*.

## Approved For Use On:

Sheep

## Active Ingredient:

ivermectin..... 0.08%

## Product Advantages:

- **Unique chemistry-unrelated to existing formulations.** Because Ivermectin Sheep Drench is chemically different, cross-resistance is unlikely in parasite strains which may have developed resistance to other wormers.
- **Broad-spectrum effectiveness:** One low-volume dose effectively treats and controls many gastrointestinal roundworms (including *Haemonchus contortus*), lungworms, and nasal bots.
- **Easy administration:** Ivermectin Sheep Drench can be administered with any standard drenching equipment or any equipment that provides a consistent dose volume.



## Direction For Use:

Ivermectin Sheep Drench may be used in any standard drenching equipment or in any equipment which provides a consistent dose volume. Ivermectin Sheep Drench is administered orally at a dose of 3.0 mL (2.4 mg ivermectin) per 26 lbs body weight or 200 mcg ivermectin per kilogram of body weight.

Coughing may be observed in some animals during and for several minutes following drenching.

## Packaging:

240mL, 6 bottles per case, UPC # 7-45801-11067-0  
· contains 20 - 100 lb. doses

960 mL, 4 bottles per case, UPC # 7-45801-11068-7  
· contains 83 - 100 lb. doses



Ivermectin Sheep Drench

062122

# Ivermectin Sheep Drench (ivermectin)

NDC 30798-637-12

0.08% Solution  
Parasiticide

NDC 30798-637-16



## Ivermectin SHEEP DRENCH

(ivermectin)  
0.08% Solution  
Parasiticide



**For the Treatment and Control  
of Worms and Bots of Sheep**

**83-100 lb Doses**

See booklet for additional information.  
**Restricted Drug** (California) - Use Only as Directed  
**For Animal Use Only - Keep Out of Reach of Children**  
**Net Contents: 32.46 fl oz (1 qt 0.46 fl oz) 960 mL**  
Approved by FDA under ANADA # 200-327

Manufactured For:  
DURVET, INC. Blue Springs, MO U.S.A. [www.durvet.com](http://www.durvet.com)

## Ivermectin Sheep Drench 0.08% Solution (ivermectin)

Approved by FDA under ANADA # 200-237

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

**FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE.**

**PRODUCT DESCRIPTION:** Ivermectin Sheep Drench is a ready-to-use, free-flowing solution of ivermectin. It is formulated to deliver the recommended dose rate of 0.2 mg ivermectin per 1 kg body weight given orally at a volume of 3.0 mL per 26 lbs body weight.

**INDICATIONS:** Ivermectin Sheep Drench provides treatment and control of adult and fourth-stage larvae of the following parasites: Gastrointestinal Roundworms – *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia curticiei*, *Nematodirus spathiger*, *N. battus*, and *Oesophagostomum columbianum*; Lungworms – *Dictyocaulus filaria*; and all the larval stages of Nasal Bot – *Oestrus ovis*. It also provides treatment and control of adult forms only of the following Gastrointestinal Roundworms – *Haemonchus placei*, *Cooperia oncophora*, *Strongyloides papillosus*, *Oesophagostomum venulosum*, *Trichuris ovis*, and *Chabertia ovina*.

1

**DOSAGE AND ADMINISTRATION:** Ivermectin Sheep Drench may be used in any standard drenching equipment or in any equipment which provides a consistent dose volume.

Ivermectin Sheep Drench is administered orally at a dose of 3.0 mL (2.4 mg ivermectin) per 26 lbs body weight or 200 mcg ivermectin per kilogram of body weight.

Coughing may be observed in some animals during and for several minutes following drenching.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

**RESIDUE WARNING:** Do not treat sheep within 11 days of slaughter.

The Safety Data Sheet (SDS) contains more detailed occupational safety information.

To report adverse effects in users, to obtain an SDS, or for assistance call 1-800-650-4899.

**PRECAUTIONS:** Ivermectin (ivermectin) Sheep Drench has been formulated for use in sheep **only**. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

**OTHER WARNINGS:** Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.

2

Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.

Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method).

A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

**Keep this and all drugs out of reach of children.**

Refrain from smoking and eating when handling. Avoid contact with eyes. Immediately wash hands and any spills on the skin with plenty of soap and water following use.

**Restricted Drug** (California) - Use Only as Directed.

**Environmental Safety:** Studies indicate that when ivermectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time. Free ivermectin may adversely affect fish and certain water-born organisms on which they feed. Do not permit water runoff from feedlots to enter lakes, streams, or ground water. Do not contaminate water by direct application or by the improper disposal of drug containers. Spills should be contained and soaked up with absorbent towels or

3

  
[www.durvet.com](http://www.durvet.com)

062122

## Ivermectin Sheep Drench

# Ivermectin Sheep Drench (ivermectin)

NDC 30798-637-12

0.08% Solution  
Parasiticide

into loose soil. Gloves should be worn to prevent skin exposure. All the collected materials (contaminated towels and soil), as well as all empty drug containers, should be placed in an impervious film (plastic) bag and disposed of by incineration or in an approved landfill.

**STORAGE INFORMATION:**

Store at 68° - 77° F (20° - 25° C). Excursions between 59° - 86° F (15° - 30° C) are permitted.

Approved by FDA under ANADA # 200-327

ISS22XB04

4



5



062122

Ivermectin Sheep Drench