

# PURINA® PRECON® COMPLETE

**Specie:** Beef**Type of Feed:** Complete**Form of Feed:** Pellet

**General Description:** Precon® Complete is a complete pelleted ration for preconditioning calves on farm or receiving calves in backgrounding operations or feedlots.

## PURINA® PRECON® COMPLETE

COMPLETE FEED FOR STARTER FEEDLOT CATTLE

**CAUTION: USE ONLY AS DIRECTED**

### GUARANTEED ANALYSIS

Crude Protein, (Min) .....	12.00 %
(This includes not more than 2.00% equivalent crude protein from non-protein nitrogen).	
Crude Fat, (Min) .....	1.00 %
Crude Fiber, (Max) .....	26.00 %
Calcium (Ca), (Min) .....	0.60 %
Phosphorus (P), (Min) .....	0.40 %
Salt (NaCl), (Min) .....	0.48 %
Potassium (K), (Min) .....	1.10 %
Vitamin A, I.U./lb, (Min) .....	5000

### NON-MEDICATED FEEDING DIRECTIONS:

Feed as a complete ration at the rate of 10 to 15 pounds per head per day to: pre-conditioned beef cattle and pre-weaned and weaned beef calves or to: beef calves upon arrival in a feed yard for a period of 7 to 10 days.

### CAUTION:

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or infested feed to animals as it may cause illness, performance loss or death.

### IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices: 1. Do not allow animals to run out of feed. 2. When making a ration change, allow 7-10 days for animals to adjust to the new ration. 3. Provide adequate feeder space for each animal. Feeders should be well protected and well managed to prevent feed from becoming wet and molding. Do not allow fines to accumulate in feed trough. Never allow the feeder to run empty or to develop flow problems. 4. Fermented feeds (high moisture grains and silages) should be limited to 10% of the diet for young or stressed cattle. 6. Stressed cattle should be placed in clean, uncrowded environments. 7. Provide salt free choice. 8. Provide fresh, clean water near the feeding area. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumptions and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

### Available Options:

Product	Product No.	Active Drug Ingredient(s)
Precon® Complete	0004686 – Bulk; 0004689 - Bag	None
*Precon® Complete LW AS87.5	0001098 – Bulk; 0001099 - Bag	Chlortetracycline, Sulfamethazine
*Precon® Complete DX.00187	0004745 – Bulk; 0004747 - Bag	Decoquinatate
*Precon® Complete B30 AU70	3003055 (-201 Bulk; -206 Bag)	Chlortetracycline, Lasalocid
*Precon® Complete AU70	0031755 – Bulk; 0017833 - Bag	Chlortetracycline
*Precon® Complete AS70	0001100 – Bulk; 0001102 - Bag	Chlortetracycline, Sulfamethazine

\*Must be fed according to feed label. Please see Purina® representative for information on individual product options and availability.