

Disease Protection



Different

- Protects against as many as 7 diseases in 1 vaccine
- Fewer needle sticks mean less pain for the horse
- Multiple combinations of vaccines packaged in pre-filled syringes
- · Peel-off sticker for easy recordkeeping

Dependable

- Safety studies of 880 horses showed 98% were reaction-free after two doses with Equi-Jec® 7(1)
- Safe for pregnant mares
- Safe for horses 4 months of age or older. Equi-Jec® 2 is safe for horses 6 months of age or older.
- Long duration of immunity: 12 months for West Nile Virus



		Sleeping Sickness				vaccinate video: 国際程序			
Coo	Product	Eastern Encephalomyelitis (EEE)	Western Encephalomyelitis (WEE)	Venezuelan Encephalomyelitis (VEE)	West Nile Virus (WNV)	Tetanus (T)	Influenza** (Flu)	Herpesvirus** (Rhino)	
	Equi-Jec® 7	•	•	•	•	•	•	•	
	Equi-Jec® 6	•	•		•	•	•	•	
	Equi-Jec° 5	•	•			•	•	•	
	Equi-Jec° 2						•	•	
	Equi-Jec® WNV+EWT	•	•		•	•			
	Equi-Jec° WNV								

(1) data on file

**See individual product label for specifics of each product.


Encephalomyelitis-Rhinopneumonitis-Influenza-West Nile Virus Vaccine

Tetanus Toxoid

Equi-Jec® 7

For Use In Animals Only

For Use In Animals Only Indications: For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by eastern, western, and venezuelan encephalomyelitis and tetanus; as an aid in the reduction of respiratory disease caused by equine influenza virus (EIV) (product contains Kentucky Lineage (KY/95), Florida sublineage clade 1 (OH/03), and Eurasian Newmarket/2/93 (NM 2/93) EIV strains), and equine herpesvirus type 1 (EHV-1) and type 4 (EHV-4); as an aid in the reduction of severity of clinical disease, and an aid in the prevention of viremia and mortality caused by west nile virus (WNV). Safe for use in pregnant mares. The duration of immunity (DOI) is at least 6 months for OH/03 and at least 12 months for WNV.



1 Dose/2 mL

Equine Rhinopneumonitis-Influenza Vaccine

Killed Virus

Equi-lec® 2

For Use In Animals Only

Indications: For vaccination of healthy horses 6 months of age or older as an aid in the reduction of respiratory disease age or older as an aid in the reduction of respiratory diseas caused by equine influenza virus (EIV) types A2 North American and A2 Eurasian (product contains Kentucky Lineage (KY/95), Florida sublineage clade 1 (OH/03), and Eurasian Newmarket/2/93 (NM 2/93) EIV strains), and equine herpesvirus type 1 (EHV-1) and type 4 (EHV-4). Safe for use in pregnant mares.



Encephalomyelitis-Rhinopneumonitis-Influenza-West Nile Virus Vaccine

Eastern & Western, Killed Virus, Tetanus Toxoid

Equi-lec® 6

For Use In Animals Only

Indications: For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by eastern and western encephalomyelitis and tetanus; as an aid in the reduction of respiratory disease caused by equine influenza virus (EIV) (product contains Kentucky Lineage (KY/95), Florida sublineage clade 1 (OH/03), and Eurasian Newmarket/2/93 (NM 2/93) EIV strains), and equine herpesvirus type 1 (EHV-1) and type 4 (EHV-4); as an aid in the reduction of severity of clinical disease, and an aid in the reduction or severity of clinical disease, and an aid in the prevention of viremia and mortality caused by west nile virus (WNV), Safe for use in pregnant mares. The duration of immunity (DOI) is at least 6 months for OH/03 and at least 12 months for WNV.



1 Dose/1 mL

Encephalomyelitis-West Nile Virus Vaccine

Eastern & Western, Killed Virus

Tetanus Toxoid

Equi-lec® WNV + EWT

For Use In Animals Only

Indications: For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by eastern and western encephalomyelitis and tetanus; and as an aid in the reduction of severity of clinical disease, and an aid in the prevention of viremia and mortality caused by west nile virus (WNV). Safe for use in pregnant mares. The duration of immunity (DOI) is at least 12 months for WNV.



1 Dose/1 mL

Encephalomyelitis-Rhinopneumonitis-Influenza Vaccine


Eastern & Western, Killed Virus

Tetanus Toxoid

Equi-Jec® 5

For Use In Animals Only

Indications: For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by eastern and western encephalomyelitis and tetanus; and as an aid in the reduction of respiratory disease caused by equine influenza virus (EIV) (product contains Kentucky Lineage (KY/95), Florida sublineage clade 1 (OH/03), and Eurasian Newmarket/2/93 (NM 2/93) EIV strains), and equine herpesvirus type 1 (EHV-1) and type 4 (EHV-4). Safe for use in pregnant mares. The duration of immunity (DOI) is at least 6 months for OH/03.



1 Dose/1 mL

West Nile Virus Vaccine

Equi-Jec® WNV

For Use In Animals Only

Indications: For vaccination of healthy horses 4 months of age or older as an aid in the reduction of severity of clinical disease, and an aid in the prevention of viremia and mortality caused by west nile virus (WNV). Safe for use in pregnant mares. The duration of immunity (DOI) is at least 12 months for WNV.

