

SweetPro®

(114 kg.) **Net Wt. 250 lbs.**

Super 25
POOR FORAGE / GUT HEALTH

Guaranteed Analysis

Crude Protein	Minimum 25.00%
(includes not more than 8% equivalent protein from non-protein-nitrogen)	
Crude Fat	Minimum 5.00%
Crude Fiber	Maximum 5.00%
Calcium (Ca)	Minimum 2.75% - Maximum 3.25%
Phosphorus (P)	Minimum 1.70%
Salt (NaCl)	Minimum 12.00% - Maximum 13.50%
Magnesium (Mg)	Minimum 3.25%
Potassium (K)	Minimum 1.00%
Sulfur (S)	Minimum 1.50%
Copper (Cu)	Minimum 600 ppm
Selenium (Se)	Minimum 5.0 ppm - Maximum 5.25 ppm
Zinc (Zn)	Minimum 1,200 ppm
Manganese (Mn)	Minimum 750 ppm
Cobalt (Co)	Minimum 4.0 ppm
Iodine (I)	Minimum 8.0 ppm
Vitamin A	Minimum 100,000 IU/lb.
Vitamin D	Minimum 20,000 IU/lb.
Vitamin E	Minimum 80 IU/lb.
Alpha-Amylase	Minimum 3,393 ug starch hydrolyzed/min/oz.
Amyloglucosidase	Minimum 513 ug maltose hydrolyzed/min/oz.

Ingredients: Corn Condensed Distillers Solubles, Corn Distillers Dried Grains with Solubles, Salt, Fermented Corn Protein, Magnesium Oxide, Mono-Calcium Phosphate, Ammonium Sulfate, Calcium Carbonate, Corn Gluten Meal, Yeast Culture, Biuret, Vegetable Oil (Flax and Canola Oil), Calcium Sulfate, Potassium Chloride, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Super 25

SweetPro® Super 25 lick blocks are designed for the essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Super 25 should be used for cattle on below average to poor forages. Daily intake will vary depending on forage quality and nutritional needs.



Feeding Instructions:

1. Before feeding SweetPro® Super 25 lick blocks for the first time, animals should be well fed.
2. Provide access for animals to have plenty of clean, fresh water at all times.
3. Have one lick block available per 25 head of beef cows.
4. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
5. Place lick blocks near water sources or areas cattle frequent regularly.
6. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
7. Other salt and minerals may be provided free choice to balance specific needs.
8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs. per day.
9. Once animals consistently consume more than 1.25 lbs/hd/day, transition to SweetPro Magnum. If daily consumption is consistently below .75 lbs/hd/day, transition to SweetPro FiberMate20.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

CAUTION: USE AS DIRECTED

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**

Do not use in association with another feed containing supplemental selenium.

Control #: _____



Guaranteed by Harvest Fuel, Inc., Walhalla, ND / Mfd. by SweetPro Feeds, Horton, KS, 66439 / 800.327.9222

All rights reserved. SweetPro is a registered trademark of Harvest Fuel, Inc. **Made in U.S.A.** Printed with soy inks.



8 91469 00032 4